

**Dangerous and Error Prone Abbreviations PC.A10.2**

Abbreviations	Intended Meaning	Misinterpretation	Correction
µg	Microgram	Mistaken as “mg”	Use “mcg”
AD, AS, AU	Right ear, left ear, each ear	Mistaken as OD, OS, OU (right eye, left eye, each eye)	Use “right ear,” “left ear,” or “each ear”
OD, OS, OU	Right eye, left eye, each eye	Mistaken as AD, AS, AU (right ear, left ear, each ear)	Use “right eye,” “left eye,” or “each eye”
BT	Bedtime	Mistaken as “BID” (twice daily)	Use “bedtime”
cc	Cubic centimeters	Mistaken as “u” (units)	Use “mL”
D/C	Discharge or discontinue	Premature discontinuation of medications if D/C (intended to mean “discharge”) has been misinterpreted as “discontinued” when followed by a list of discharge medications	Use “discharge” and “discontinue”
IJ	Injection	Mistaken as “IV” or “intrajugular”	Use “injection”
IN	Intranasal	Mistaken as “IM” or “IV”	Use “intranasal” or “NAS”
HS	Half-strength	Mistaken as bedtime	Use “half-strength” or “bedtime”
hs	At bedtime, hours of sleep	Mistaken as half-strength	Use “half-strength” or “bedtime”
IU**	International unit	Mistaken as IV (intravenous) or 10 (ten)	Use “units”
o.d. or OD	Once daily	Mistaken as “right eye” (OD-oculus dexter), leading to oral liquid medications administered in the eye	Use “daily”
OJ	Orange juice	Mistaken as OD or OS (right or left eye); drugs meant to be diluted in orange juice may be given in the eye	Use "orange juice"

## Lifesong Hospice and Palliative Care

### Patient Care Policies and Procedures

Abbreviations	Intended Meaning	Misinterpretation	Correction
Per os	By mouth, orally	The “os” can be mistaken as “left eye” (OS-oculus sinister)	Use “PO,” “by mouth,” or “orally”
q.d. or QD**	Every day	Mistaken as q.i.d., especially if the period after the “q” or the tail of the “q” is misunderstood as an “i”	Use “daily”
qhs	Nightly at bedtime	Mistaken as “qhr” or every hour	Use “nightly”
qn	Nightly or at bedtime	Mistaken as “qh” (every hour)	Use “nightly” or “at bedtime”
q.o.d. or QOD**	Every other day	Mistaken as “q.d.” (daily) or “q.i.d. (four times daily) if the “o” is poorly written	Use “every other day”
q1d	Daily	Mistaken as q.i.d. (four times daily)	Use “daily”
q6PM, etc	Every evening at 6 PM	Mistaken as every 6 hours	Use “daily at 6 PM” or “6 PM daily”
SC, SQ, sub q	Subcutaneous	SC mistaken as SL (sublingual); SQ mistaken as “5 every;” the “q” in “sub q” has been mistaken as “every” (e.g., a heparin dose ordered “sub q 2 hours before surgery” misunderstood as every 2 hours before surgery)	Use “subcut” or “subcutaneously”
ss	Sliding scale (insulin) or ½ (apothecary)	Mistaken as “55”	Spell out “sliding scale;” use “one-half” or “½”
SSRI	Sliding scale regular insulin	Mistaken as selective-serotonin reuptake inhibitor	Spell out “sliding scale (insulin)”
SSI	Sliding scale insulin	Mistaken as Strong Solution of Iodine (Lugol's)	Spell out “sliding scale (insulin)”
i/d	One daily	Mistaken as “tid”	Use “1 daily”
TIW or tiw	3 times a week	Mistaken as “3 times a day” or “twice in a week”	Use “3 times weekly”
U or u**	Unit	Mistaken as the number 0 or 4, causing a 10-fold overdose or greater (e.g., 4U seen as	Use “unit”

## Lifesong Hospice and Palliative Care

### Patient Care Policies and Procedures

Abbreviations	Intended Meaning	Misinterpretation	Correction
		"40" or 4u seen as "44"); mistaken as "cc" so dose given in volume instead of units (e.g., 4u seen as 4cc)	
UD	As directed ("ut dictum")	Mistaken as unit dose (e.g., diltiazem 125 mg IV infusion "UD" misinterpreted as meaning to give the entire infusion as a unit [bolus] dose)	Use "as directed"

### *Customization Suggestions: Abbreviations*

*This list is adapted from <http://www.ismp.org/Tools/errorproneabbreviations.pdf>*

*Please update at least annually*

*In addition, information from the Joint Commission on "dangerous abbreviations" is available at [https://www.jointcommission.org/facts\\_about\\_do\\_not\\_use\\_list/](https://www.jointcommission.org/facts_about_do_not_use_list/)*